


Pediatric Cardiac Arrest Algorithm



CPR Quality

- Push hard (≥1/3 of anteroposterior diameter of chest) and fast (100-120/min) and allow complete chest recoil
- Minimize interruptions in compressions
- Change compressor every 2 minutes, or sooner if fatigued
- If no advanced airway, 15:2 compression-ventilation ratio
- If advanced airway, provide continuous compressions and give a breath every 2-3 seconds

Shock Energy for Defibrillation

- First shock 2 J/kg
- Second shock 4 J/kg
- Subsequent shocks ≥4 J/kg, maximum 10 J/kg or adult dose

Drug Therapy

- **Epinephrine IV/IO dose:** 0.01 mg/kg (0.1 mL/kg of the 0.1 mg/mL concentration). Max dose 1 mg. Repeat every 3-5 minutes. If no IV/IO access, may give endotracheal dose: 0.1 mg/kg (0.1 mL/kg of the 1 mg/mL concentration).
- **Amiodarone IV/IO dose:** 5 mg/kg bolus during cardiac arrest. May repeat up to 3 total doses for refractory VF/pulseless VT or
- **Lidocaine IV/IO dose:** Initial: 1 mg/kg loading dose

Advanced Airway

- Endotracheal intubation or supraglottic advanced airway
- Waveform capnography or capnometry to confirm and monitor ET tube placement

Reversible Causes

- Hypovolemia
- Hypoxia
- Hydrogen ion (acidosis)
- Hypoglycemia
- Hypo-/hyperkalemia
- Hypothermia
- Tension pneumothorax
- Tamponade, cardiac
- Toxins
- Thrombosis, pulmonary
- Thrombosis, coronary